Te Marae Ora Ministry of Health Indicator Matrix Jul 2020 (129)

Short term <12 months

Medium term >1-5 years

Long term >10 years

Long term >10 years	DIOK FACTORS (SA)	OFFINIOR OF VERY OF VERY	LIEALTH OVOTERS (A4)
HEALTH STATUS (34)	RISK FACTORS (31) Nutrition (7)	SERVICE COVERAGE (33)	HEALTH SYSTEMS (31) Quality and safety of care (13)
Mortality by age and sex (8)	Nutrition (7)	Reproductive maternal, new-born, child and adolescent (6)	Quality and safety of care (13)
Life expectancy at birth	Exclusive breastfeeding rate 0-5months of age	Demand for family planning satisfied with modern methods	Perioperative mortality rate
Crude death rate	Incidence of low birth weight among new-borns	Contraceptive prevalence rate	Institutional maternal mortality ratio
Adolescent mortality rate Adult mortality rate between 15 and 60 years	Children under 5 years who are overweight Anaemia prevalence in children	Antenatal care coverage Births attended by skilled health personnel	Maternal death reviews ART retention rate
of age			
Under-five mortality rate	Anaemia prevalence in women of reproductive age	Postpartum care coverage - women	TB treatment success
Infant mortality rate	Children under 5 years who are stunted	Postnatal care coverage - new-born	Clinical protocols and guidelines for all specialty areas
Neonatal mortality rate Stillbirth rate	Children under 5 years who are wasted Environmental risk factors (5)	Immunisation (2) Immunisation coverage rate by vaccine for each	Availability of essential medicines and commodities Rate of adverse events among specialty areas
Mortality by course (40)	Deputation using anfalt managed dripting water	vaccine in the national schedule <5 years	Complication rate among long term care nations nanulation
Mortality by cause (10)	Population using safely managed drinking water services	Availability of vaccines against human papillomavirus, according to national programmes	Complication rate among long term care patient population
Suicide rate	Population using safely managed sanitation	and policies HIV (4)	Number and proportion of domestic patient referrals
Dark and during the said traffic injuries	services		
Death rate due to road traffic injuries	Population with handwashing facility with soap and water	People living with HIV who know their status	Number and proportion of international patient referrals
Mortality rate due to homicide	Number and proportion of reported foodborne illnesses	Prevalence of mother-to-child transmission	Ambulatory Sensitive Hospitalisation (ASH) rates for 0-4 year olds
Maternal mortality ratio	Number and proportion of reported waterborne	Antiretroviral therapy (ART) coverage	% of compliance with Package of Essential NCD
TB mortality rate	illnesses Population with primary reliance on clean fuels and	HIV viral load suppression	interventions (PEN) guidelines Utilisation and access (11)
·	technologies		
AIDS-related mortality rate Premature noncommunicable disease (NCD)	Non communicable diseases (14) Insufficient physical activity in adults	Coverage of treatment for latent TB infection (LTB)	Access to primary health care Access to palliative care
mortality			
Unconditional probability of dying between ages of 30 and 70 from cardiovascular	Insufficient physical activity in adolescents	HIV test results for TB patients	Proportion of patients who have seen a primary care provider/GP within 7 days of discharge
diseases, cancer, diabetes or chronic	Total alashal and angita (and 45, under)	LIN/iti	N
Mortality from unsafe water, unsafe sanitation and lack of hygiene	consumption	HIV-positive new and relapse TB patients on ART during TB treatment	Number and proportion of outpatient consultations
Mortality from unintentional poisoning	Age-standardised prevalence of heavy episodic drinking among adolescents and adults, as	Tuberculosis (3)	Number and proportion of consultations for oral health services
	appropriate, within the national context		SCIVILES
Fertility (3)	Alcohol-related morbidity and mortality among adolescents and adults, as appropriate, within the	Drug susceptibility testing coverage for TB patients	Number and proportion of people aged two and over who had a dental visit within the last 12 months
Crude birth rate	national context Tobacco use among persons aged 15+ years	TB treatment coverage	Inpatient admissions
Adolescent birth rate	Raised blood pressure among adults	Treatment coverage for drug-resistant TB	30-day readmission rate after hospital discharge
Total fertility rate Morbidity (13)	Raised blood glucose/diabetes in adults Raised blood glucose/diabetes in adolescents	Vector borne diseases (4) Intermittent preventive therapy for vector borne	Surgical volume Health facility density and distribution
New cases of vaccine-preventable diseases	· .	diseases during pregnancy Use of insecticide treated nets	
ivew cases of vaccine-preventable diseases	Age-standardised mean proportion of total energy intake from saturated fatty acids in person aged	Use of insecticide freated field	Hospital bed density
New cases of IHR-notifiable diseases and	18+ years Age-standardised prevalence of raised total	Treatment of confirmed dengue cases	Health workforce (1)
other notifiable diseases	cholesterol among persons aged 18+ years	noament et centiminea aongae cacce	Treath Worklove (1)
	(defined as total cholesterol >5mmol/L) and mean total cholesterol concentration		
NCD morbidity rate	Age-standardised prevalence of persons	Indoor residual spraying coverage for dengue	Health worker density and distribution
	consuming less than 5 total servings (400grams) of fruits and vegetables per day		
HIV incidence rate Hepatitis B incidence	Salt intake Overweight and obesity in adults	Screening and preventive care (3)	Health information (2) Birth registration
Sexually transmitted infections (STIs)	Injuries/harmful traditional practices (5)	Breast cancer screening Cervical cancer screening	Death registration
incidence rate Congenital syphilis rate	Intimate partner violence number and prevalence	Proportion of women between the ages of 30-49	Health security (1)
3		screened for cervical cancer at least once, or more	
		often, and for lower or higher age groups according to national programmes or policies	
TB incidence rate	Non-partner sexual violence number and prevalence	Mental health (2)	International Health Regulations (IHR) core capacity index
TB notification rate	Sexual violence against children (number and	Number and proportion of mental health disorders	Health financing (3)
Cancer incidence, by type of cancer	prevalence) Frequency rates of occupation injuries	Coverage of services for severe mental health	Health Care Expenditure (HCE) as a percentage of gross
HIV prevalence rate	% of injury related fatalities across all patient	disorders NCD (4)	domestic product (GDP) HCE per capita
	groups		
Hepatitis B surface antigen prevalence		% of diabetes patients receiving eye care visits/treatment from a specialist within one year	HCE as a percentage of Govt expenditure
NCD prevalence rate		Admission rates for conditions that are sensitive to outpatient (ambulatory) care delivery	
		Policies to reduce the impact on children of	
		marketing of foods and non-alcoholic beverages high in saturated fats, trans fatty acids, free sugars,	
		or salt	
		Adoption of national policies that limit saturated fatty acids and virtually eliminate partially hydrogenated	
		vegetable oils in the food supply, as appropriate,	
		within the national context and national programmes	
		Substance abuse (1)	
		Treatment coverage for alcohol and drug dependence	
		Health Specialist Visits (1) Number and proportion of population accessing	
		health services by area of speciality	